



## NCSTN gene

nicastrin

### Normal Function

The *NCSTN* gene provides instructions for making a protein called nicastrin. This protein is one part (subunit) of a complex called gamma- ( $\gamma$ -) secretase. Nicastrin plays a critical role in the assembly and stability of this complex.

The  $\gamma$ -secretase complex is located in the membrane that surrounds cells, where it cuts apart (cleaves) many different proteins that span the cell membrane (transmembrane proteins). This cleavage is an important step in several chemical signaling pathways that transmit signals from outside the cell into the nucleus. One of these pathways, known as Notch signaling, is essential for the normal maturation and division of hair follicle cells and other types of skin cells. Notch signaling is also involved in normal immune system function.

### Health Conditions Related to Genetic Changes

#### hidradenitis suppurativa


At least 11 mutations in the *NCSTN* gene have been found to cause hidradenitis suppurativa, a chronic skin disease characterized by recurrent boil-like lumps (nodules) under the skin that develop in hair follicles. The nodules tend to become inflamed and painful, and they produce significant scarring as they heal.

*NCSTN* gene mutations reduce the amount of functional nicastrin produced in cells, so less of this protein is available to act as part of the  $\gamma$ -secretase complex. The resulting shortage of normal  $\gamma$ -secretase impairs cell signaling pathways, including Notch signaling. Although little is known about the mechanism, studies suggest that abnormal Notch signaling may promote the development of recurrent nodules in hair follicles and trigger inflammation in the skin.

## Chromosomal Location

Cytogenetic Location: 1q23.2, which is the long (q) arm of chromosome 1 at position 23.2

Molecular Location: base pairs 160,343,056 to 160,358,952 on chromosome 1 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

## Other Names for This Gene

- anterior pharynx-defective 2
- APH2
- ATAG1874
- KIAA0253
- NICA\_HUMAN
- nicastrin precursor
- RP11-517F10.1

## Additional Information & Resources

### Educational Resources

- Marie Curie Bioscience Database: Notch Signaling and the Developing Hair Follicle  
<https://www.ncbi.nlm.nih.gov/books/NBK45997/>

### Scientific Articles on PubMed

- PubMed  
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28NCSTN%5BTIAB%5D%29+OR+%28nicastrin%5BTIAB%5D%29%29+AND+%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1440+days%22%5Bdp%5D>

## OMIM

- NICASTRIN  
<http://omim.org/entry/605254>

## Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology  
[http://atlasgeneticsoncology.org/Genes/GC\\_NCSTN.html](http://atlasgeneticsoncology.org/Genes/GC_NCSTN.html)
- ClinVar  
<https://www.ncbi.nlm.nih.gov/clinvar?term=NCSTN%5Bgene%5D>
- HGNC Gene Symbol Report  
[http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?q=data/hgnc\\_data.php&hgnc\\_id=17091](http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=17091)
- NCBI Gene  
<https://www.ncbi.nlm.nih.gov/gene/23385>
- UniProt  
<http://www.uniprot.org/uniprot/Q92542>

## **Sources for This Summary**

- Liu Y, Gao M, Lv YM, Yang X, Ren YQ, Jiang T, Zhang X, Guo BR, Li M, Zhang Q, Zhang P, Zhou FS, Chen G, Yin XY, Zuo XB, Sun LD, Zheng XD, Zhang SM, Liu JJ, Zhou Y, Li YR, Wang J, Wang J, Yang HM, Yang S, Li RQ, Zhang XJ. Confirmation by exome sequencing of the pathogenic role of NCSTN mutations in acne inversa (hidradenitis suppurativa). *J Invest Dermatol.* 2011 Jul;131(7):1570-2. doi: 10.1038/jid.2011.62. Epub 2011 Mar 24.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/21430701>
- Melnik BC, Plewig G. Impaired Notch signalling: the unifying mechanism explaining the pathogenesis of hidradenitis suppurativa (acne inversa). *Br J Dermatol.* 2013 Apr;168(4):876-8. doi: 10.1111/bjd.12068. Epub 2013 Jan 31.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/23020871>
- Pink AE, Simpson MA, Brice GW, Smith CH, Desai N, Mortimer PS, Barker JN, Trembath RC. PSENEN and NCSTN mutations in familial hidradenitis suppurativa (Acne Inversa). *J Invest Dermatol.* 2011 Jul;131(7):1568-70. doi: 10.1038/jid.2011.42. Epub 2011 Mar 17.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/21412258>
- Pink AE, Simpson MA, Desai N, Dafou D, Hills A, Mortimer P, Smith CH, Trembath RC, Barker JN. Mutations in the  $\gamma$ -secretase genes NCSTN, PSENEN, and PSEN1 underlie rare forms of hidradenitis suppurativa (acne inversa). *J Invest Dermatol.* 2012 Oct;132(10):2459-61. doi: 10.1038/jid.2012.162. Epub 2012 May 24.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/22622421>

- Pink AE, Simpson MA, Desai N, Trembath RC, Barker JN.  $\gamma$ -Secretase mutations in hidradenitis suppurativa: new insights into disease pathogenesis. *J Invest Dermatol.* 2013 Mar;133(3):601-7. doi: 10.1038/jid.2012.372. Epub 2012 Oct 25. Review.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/23096707>
- Wang B, Yang W, Wen W, Sun J, Su B, Liu B, Ma D, Lv D, Wen Y, Qu T, Chen M, Sun M, Shen Y, Zhang X. Gamma-secretase gene mutations in familial acne inversa. *Science.* 2010 Nov 19; 330(6007):1065. doi: 10.1126/science.1196284. Epub 2010 Oct 7.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/20929727>

---

Reprinted from Genetics Home Reference:

<https://ghr.nlm.nih.gov/gene/NCSTN>

Reviewed: December 2013

Published: March 21, 2017

Lister Hill National Center for Biomedical Communications

U.S. National Library of Medicine

National Institutes of Health

Department of Health & Human Services